


TARRANT COUNTY

# Arboviral Surveillance and Mosquito Control Program

---

Tarrant County Public Health

2/12/2020



Disclaimer: Vector surveillance and control is a dynamic process and as such this document is subject to change during the operational period.

Rev. 2/2019

## Table of Contents

I.	Introduction	3
A.	West Nile virus and St. Louis encephalitis virus	3
B.	Chikungunya virus, Zika virus, and Dengue virus	4
II.	Tarrant County management area description	5
III.	Surveillance for arboviruses and arboviral diseases	5
A.	West Nile virus and St. Louis encephalitis virus surveillance in mosquitoes	5
ii.	Off-season surveillance	7
B.	Dengue virus, Zika virus, and chikungunya virus surveillance in mosquitoes and surveillance for Aedes mosquitoes	7
i.	Mosquito surveillance and control near imported cases	7
C.	Human surveillance:	9
D.	Birds and sentinel chicken surveillance (WNV/SLEV only):	9
E.	Equine surveillance (WNV/SLEV Only):	9
IV.	Prevention and control	10
A.	Larval mosquitoes	10
B.	Adult mosquitoes	11
C.	Resistance management	12
V.	Education, outreach, and media	12
VI.	Mosquito complaints	13
	Appendix A: Trap Location	14
	Appendix B: Internal notification of positive mosquito pools and trap location	15
	Appendix C: List of documents, emergency response, and other documentation	16
	General Documents	16
	Emergency response	16
	Personal protective equipment (PPE)	16
	Spill kit	17
	Other documents	17
	Appendix D: Forms Utilized in Zika Response	18
		18
	Appendix E: Larvicides	22

Microbial	22
Bacillus thurigiensis isreaeensis (Bti)	22
Lysinibacillus sphaericus formerly Bacillus sphaericus (Bs)	22
Spinosad	22
Chemical/Physical	22
Insect Growth Regulators (IGRs)	22
Monomolecular Films (MMF)	23
Oils	23
Appendix F: <i>Gambusia affinis</i>	24
Appendix G: Adulticides	26
Chemical	26
Pyrethroids	26
Organophosphates	26
Appendix H. Phased response Guidelines	27
Table 1 Tarrant County Public Health Guidelines for Phased Response to WNV/SLEV Surveillance	27
Table 2 Tarrant County Public Health Guidelines for Phased Response to human cases of chikungunya, dengue and Zika	29
Appendix I: Laws pertaining to vector control	32
Texas Health and Safety Code	32
Title 5, Subtitle A, Chapter 341, Subchapter B	32
Title 5, Subtitle A, Chapter 343, Subchapter B	32
Title 5, Subtitle A, Chapter 341, Subchapter C	32
Glossary of acronyms	33
Works cited	35

## List of Tables and Figures

Figure 1. Locations of in-season and off-season gravid trap locations in Tarrant County. ....	14
Figure 2. Aerial picture of live wells for storage of <i>Gambusia affinis</i> .....	24
Figure 3. Location of TRWD fish hatchery in northwest Tarrant County.....	25
Table 1. Tarrant County Public Health Guidelines for phased response to mosquito surveillance.....	16

## I. Introduction

Arboviral and other arthropod-borne diseases are those in which the disease-causing agent is spread by the bite of an infected arthropod. As there are many arboviral diseases that have the potential to show up in any area, at any time, we will be focusing this document on five viruses of concern in North Texas. The agents of concern currently include West Nile virus (WNV), St. Louis encephalitis virus (SLEV), chikungunya virus (CHIKV), dengue virus (DENV), and Zika virus (ZIKV). Due to the nature of these diseases, WNV and SLEV can be discussed together as zoonotic diseases, shared between animals and humans, whereas CHIKV, DENV, and ZIKV can be discussed together as epidemic diseases, shared among humans. The content of this document is subject to change dependent on the most current information and potentially upcoming events. Tarrant County Public Health (TCPH) conducts surveillance for the presence of WNV and SLEV by studying, surveying, sampling and testing local *Culex* populations in unincorporated portions of Tarrant County. The presence of CHIKV, DENV, and ZIKV, on the other hand, will be determined predominantly by the presence of these viruses in the human population; however, the presence and abundance of *Aedes aegypti* and *Aedes albopictus* will be surveyed to help determine high risk areas. In the middle of 2016 until the current date, outbreaks of Yellow Fever emerged around Africa and Brazil. Many throughout North America have not been vaccinated for this devastating illness. Tarrant County Public Health recognizes that though there is a vaccine for Yellow Fever, there is still a risk for this disease. Responses to a possible outbreak of Yellow Fever would be similar to that of CHIKV, DENV, and ZIKV with the addition of vaccine intervention.

### A. West Nile virus and St. Louis encephalitis virus

West Nile virus and St. Louis encephalitis virus are endemic mosquito-borne viruses that have the potential to cause febrile disease, myelitis, encephalitis, and/or meningitis in humans and other animals. Approximately 80% of people who acquire these viruses will not experience symptoms. Of the twenty percent who do, roughly one percent will develop serious symptoms. WNV was first isolated in a febrile patient in the African country of Uganda, circa 1937. After first appearing in the US in New York City in the summer of 1999, it spread west eventually reaching north Texas in 2002. (AAM, 2013). Both West Nile virus and St. Louis encephalitis virus are predominantly carried by the *Culex* genus of mosquito. The primary vector of these diseases in North Texas is known to be *Culex quinquefasciatus* (Ward, 2002). *Culex restuans* and *Culex nigripalpus* have also tested positive for WNV and will be considered secondary vectors. Therefore, much of this program will target surveillance and control of these disease carrying mosquitoes. Other species of *Culex* that may be vectors for these diseases will also be tested when sampled in abundance.

*Culex* surveillance will be conducted year-round. Regular in-season testing will begin the first full week of April and end in mid-November when cooler temperatures are most likely. Off-season surveillance will be conducted to monitor the mosquito populations in cooler winter months. Female *Culex* mosquitoes over-winter as adults which are virtually inactive at times of cold temperatures

(Strickman, 1988). Minimal surveillance will be conducted when temperatures remain consistently low (below 8°C). Being nocturnal, these mosquitoes are most active between dusk and dawn.

WNV and SLEV have a bird-mosquito-bird transmission cycle that includes humans, horses and other animals as dead-end hosts. This means that a mosquito can only acquire the virus from a bird and cannot acquire the virus from a human or horse (that is infected with WNV or SLEV) because of the low number of viral particles in our bloodstreams. After a period of time, the infected mosquito can pass these viruses on to other animals. Data pertaining to information about WNV/SLEV in human, horse and bird populations will also be considered as important information in decision making and recommendations for the county as a whole. After a period collecting robust data (at least 5 years), Tarrant County Public Health will assess establishing thresholds to determine mosquito control activities. These may include mosquito infection rates, vector indices, abundance of mosquitoes, weather patterns, and timing/rate of increase of mosquito positivity.

## B. Chikungunya virus, Zika virus, and Dengue virus

Chikungunya virus (CHIKV), Zika virus (ZIKV), and dengue virus (DENV) are three arboviruses that are a potential threat to people living in North America. All three viruses share the primary vector *Aedes aegypti* and a potential secondary vector *Aedes albopictus*, which are present and abundant in many areas of Tarrant County. These mosquitoes behave differently than *Culex* mosquitoes, so surveillance for these viruses will be different than that of WNV/SLEV.

Chikungunya virus is an *alphavirus* in the family *Togaviridae*. Disease manifested by this virus, known as chikungunya fever (CHIK), can result in high fever, headaches, intense joint pain and sometimes a rash. It was first isolated in 1952 on the Makonde plateau in Tanzania, Eastern Africa. The word “chikungunya” roughly translates to “that which bends up” in the Makonde dialect which describes the actions of those experiencing symptoms of the disease. Since 2004, the distribution of the virus has been spreading from the African islands, to Asia, Europe and has most recently made a stronghold in the Caribbean and other popular North American island destinations (PAHO, 2011). Chikungunya may be severely debilitating but is rarely fatal. Symptoms typically last less than two weeks, but may persist for months and in rare cases, even years.

Dengue virus (*Flaviviridae: flavivirus*) continues to spread around tropical and subtropical regions. Disease resulting from DENV can range from febrile to hemorrhagic and it exists in four different serotypes, 1, 2, 3 and 4. The symptoms of dengue fever can be similar to CHIK, however the onset of fever is more gradual, there is more commonly a rash, and it is more likely to be fatal (PAHO, 2011). Once an infected person overcomes illness from one serotype of DENV, they become susceptible to develop more severe dengue fever symptoms upon reinfection with another serotype (WHO, 2009). Recently, local transmission has been documented in the Florida Keys (Richards *et al.* 2012) and happens periodically on the Texas-Mexico border (Rodrigues-Tan and Weir, 1998).

Like dengue virus, Zika virus is in the family *Flaviviridae*. Symptoms from acquisition of this virus are also similar to CHIKV and DENV. Most cases result in a maculopapular rash (flattened raised red patches) on the torso, joint pain and conjunctivitis. Symptoms are believed to be less severe than that of CHIKV and speculated to be manifested by roughly 20% of people who acquire the virus. Guillain-Barre Syndrome and encephalitis-like diseases have been associated with Zika in a very small percentage of the people who have developed Zika-related illness. Deaths associated with Zika virus are rare. The cause of major concern with the spread of Zika is the effects it has on fetuses if women are infected during pregnancy. In April of 2016, CDC scientists announced that Zika can cause microcephaly and other severe fetal brain defects (Rasmussen *et al* 2016). Microcephaly can be defined as a condition where a baby is born with an abnormally small head, likely due to the under-development of the brain. For more information on microcephaly, visit: <https://www.cdc.gov/ncbddd/birthdefects/microcephaly.html> Pregnant women should avoid travel to places where Zika is being transmitted. Most cases of Zika-illness in the United States are related to international travel. In 2016 two outbreaks of Zika occurred in the intercontinental US in Miami, Florida and Brownsville, Texas. There were an additional 4 cases of Zika in Hidalgo and Cameron Counties in 2017. Though most cases of Zika are transmitted through the bite of an infected mosquito, it may also be transmitted sexually.

All three of these viruses are transmitted by mosquitoes in the genus *Aedes*, subgenus *Stegomyia*. These mosquitoes are unlike *Culex* mosquitoes in the fact that they are diurnal (active during the day), overwinter as eggs, and predominantly take bloodmeals from humans. Surveillance for these mosquitoes will take place during our in-season WNV/SLEV surveillance (see section A.) utilizing BG Sentinel traps baited with CO<sub>2</sub>. Surveillance for these mosquitoes will only be utilized to find potential problem areas and not for surveillance of these viruses since there is no autochthonous transmission and humans are the current preferred sentinel.

## **II. Tarrant County management area description**

Tarrant County Public Health (TCPH) is responsible for all areas of unincorporated Tarrant County. TCPH will also provide assistance and recommendations to incorporated entities within the county.

## **III. Surveillance for arboviruses and arboviral diseases**

### **A. West Nile virus and St. Louis encephalitis virus surveillance in mosquitoes**

Surveillance for WNV/SLEV in the local mosquito population enables TCPH to assess the risk for transmission to humans by calculating the minimum infection rate and the vector index. These methods are recommended for risk assessment by the Centers for Disease Control and Prevention (CDC) in their

*West Nile Virus in the United States: Guidelines for Surveillance, Prevention and Control*. Mosquitoes will typically test positive for WNV/SLEV before human cases appear.

The Environmental Health Division (EHD) of TCPH will perform trapping of adult *Culex* mosquitoes in unincorporated portions of Tarrant County (for more information about trap location, see Appendix A). Cities may collaborate with TCPH by trapping adult mosquitoes at chosen locations within their city. There are greater than 200 static trap locations around the county. Each entity will be expected to perform certain duties as stated in the following protocol:

EHD will supply traps, bins, nets, batteries and delivery containers based on population/size of city and perform maintenance on these traps during the winter season.

#### Seasonal surveillance

- A. Collaborating cities will be expected to: Brew gravid water with grass clippings/alfalfa/hay and tap or natural collected water for a minimum of 7 days (14 days preferred), set traps overnight to be collected the following morning (once/week), and deliver mosquitoes in appropriate containers and with completed paper work (including if no mosquitoes were captured) to TCPH room 2300 (vector control room) in supplied containers no later than 24 hours prior to set date to be tested. If all traps yield no mosquitoes, paperwork may be sent via e-mail to the vector control supervisor or fax) to (817) 321-4961(with ATTN: Vector Control).
- B. EHD will: Freeze mosquitoes prior to identification, identify, enumerate, and place up to 50 mosquitoes of the appropriate vector species into each tube. Mosquitoes may be stored in a -30° C freezer and delivered to North Texas Regional Laboratory (NTRL) no later than 12:00 PM on a predetermined date to be tested.
- C. NTRL will: Store frozen samples until ready for processing (normally performed on Tuesdays, Wednesdays and Thursdays), produce homogenates, extract RNA from samples, run a real-time reverse-transcriptase polymerase chain reaction (RT-PCR) assay for detection of WNV and SLEV, determine positive/negative results and report results to EHD the morning of the day following laboratory submission, barring unforeseen circumstances.
- D. EHD will: Notify municipalities of any mosquito pools that test positive for WNV or SLEV via telephone and secondarily via e-mail. All results will be received via e-mail. Vector control personnel or designee will also complete the internal notification (see Appendix B). Locations and results of WNV/SLEV trap sites will be added to the Tarrant County website by vector control personnel or designee before receiving results, but results will not be posted to the TCPH website until the day after receiving results from the lab to allow for the municipalities to properly notify their personnel/public.
- E. EHD will: Vector control personnel or designee will report positive mosquito pool results to Texas Department of State Health Services (DSHS) once/week on Fridays, barring unforeseen circumstances.

## ii. Off-season surveillance

- A. Collaborating cities will be expected to: Brew gravid water with grass clippings/alfalfa/hay and tap or natural collected water for a minimum of 7 days (14 days preferred), set traps overnight to be collected the following morning (once/every other week) barring temperatures less than 37°F (Strickman, 1988), and deliver mosquitoes and completed paper work (including if no mosquitoes were captured) to TCPH room 2300.
- B. Off-season collection will follow the same procedures as in-season trapping other than section A. An electronic calendar will be supplied to participating cities.

Off-season trapping should follow the same procedures as the in-season trapping.

## B. Dengue virus, Zika virus, and chikungunya virus surveillance in mosquitoes and surveillance for *Aedes* mosquitoes

Surveillance for CHIKV, ZIKV, and DENV in mosquitoes will only happen on a case-by-case basis where one of these viruses would be most likely to occur. This would be strictly for data collection purposes only and would not determine a response for mosquito treatment. Due to the nature of the way these viruses cycle in the environment, strictly between humans and mosquitoes, it is highly unlikely that these viruses will be found in the mosquito populations *before* human cases appear. Therefore, the dominant form of surveillance for the virus itself will be through recording confirmed human cases. Tarrant County Public Health and municipalities within Tarrant County deploy BG Sentinel traps to monitor the populations of *Aedes aegypti* and *Aedes albopictus*. Static trap locations have been chosen based on human population density and breeding habitat availability. BG Sentinel traps with the addition of dry ice should also be placed at suspect case addresses during warmer seasons (during the WNV in-season period). TCPH has distributed BG Sentinel traps for the surveillance of *Ae aegypti* and *Ae albopictus* mosquitoes to our municipalities. Results of BG Sentinel traps will be turned in with gravid trap samples and results will be returned with gravid trap results. It is important to keep these traps in static locations to monitor these mosquito populations over time.

### i. Mosquito surveillance and control near imported cases

Suspected imported cases of CHIK, Zika or dengue fever will be determined by the Division of Epidemiology at TCPH. Suspect imported cases must have a travel history that includes a country in which CHIKV, ZIKV and/or DENV are endemic, epidemic, or currently circulating. Upon notification of a suspected human case from the Division of Epidemiology, a member of the Vector Control team will work collaboratively with the appropriate municipality personnel to inspect the property of the patient

for sources of *Aedes* breeding and to determine subsequent mosquito control activities. It is anticipated that municipality personnel will be able to handle complaints independently after a few inspections. Permission forms should be utilized to enter private residences (Appendix D). Personnel conducting the inspection will record observations on the backyard mosquito checklist (Appendix E). It should be noted here that risk is determined by a number of factors. These include viremia of the patient while they were present in the county, the number of vector mosquitoes found on or near the property, the use of and type of repellents utilized during the viremic period, and/or reports of patient outdoor activity/reports of being bitten by vector mosquitoes. BG Sentinel traps should be deployed with the addition of dry ice for a twenty-four hour period during times when *Ae aegypti* and *Ae albopictus* are known to be active (typically from May to November) to abundance of adult mosquitoes. BG Sentinel trap samples should be brought to Tarrant County Vector Control Staff the day following trap placement where vector control staff may be able to make risk assessments on a case-by-case basis. Adulticiding the patient residence and all adjoining properties may be recommended where high risk is determined to be present along with larviciding and education to all residences within a 150 meter radius. Handing out repellents including *N, N, N* diethyl-*meta*-toluamide (DEET) and properly labeled larvicides within the neighborhood may also be encouraged. All larval source containers should be overturned or treated with the appropriate larvicide. Upon observation of adult mosquitoes, adulticiding should be considered. Adulticiding may include thermal fogging or residual spray to surrounding resting areas (low-laying, non-flowering plant life and eaves). Treatment of flowering plants should be avoided to minimize exposure to non-target insects. ULV targeted backpack application is not recommended in the daytime due to a lack of inversion layer where the chemical will evaporate and be ineffective. The placement of a post-treatment BG Sentinel trap is recommended to monitor effectiveness of treatment. This data should be reported to TCPH for data collection purposes. If the patient is no longer viremic and therefore cannot spread the virus to a local mosquito population, the patient will be educated about the use of repellent and encouraged to treat larval sources found on their property.

## ii. Mosquito surveillance near autochthonous cases

Autochthonous cases can be described as “locally acquired” cases. This can be determined by suspected symptoms and lack of travel to an endemic/epidemic country. Mosquito control activities on the patient’s property will be determined on the likelihood of CHIK/DENV/ZIKV fever. Once an autochthonous case is confirmed, both the Texas Department of State Health Services and the Centers for Disease Control and Prevention will be notified and consulted. Information on *Aedes* mosquito breeding habitat, prevention of mosquito bites and symptoms of these diseases near an autochthonous case will be the primary means to minimize additional cases. Mosquito control activities including source reduction, larviciding and adulticiding will ensue within a 150-meter radius around an autochthonous case, or the approximate flight range of *Aedes aegypti* and *Aedes albopictus* (PAHO, 2011). BG Sentinel traps will also be placed within a 150 meter of the human case to monitor the local mosquito population. Control measures should remain similar to that of imported cases.

### C. Human surveillance:

Arboviral diseases and exotic diseases are nationally-notifiable conditions and reportable in Texas. Most disease cases including suspected cases are reported to the Division of Epidemiology at TCPH. They are responsible for ensuring that reported human disease cases meet the Council of State & Territorial Epidemiologists (CSTE) case definitions\* and are investigated promptly. Upon confirmation of a human WNV or SLEV case, the Division of Epidemiology will notify a designee in Vector Control. In compliance with the Health Insurance Portability and Accountability Act (HIPAA), information regarding human cases will not be released on the interactive mapping website. Vector Control staff will report an intersection nearby the human case to the municipality where the case resides and TCPH recommends setting additional gravid traps near this area to test additional pools of mosquitoes for WNV/SLEV. This is recommended due to the nature of acquisition not being isolated to the patient's home and also to help protect HIPAA information in reasons to treat.

### D. Birds and sentinel chicken surveillance (WNV/SLEV only):

Passerine birds, or perching birds, serve as the main reservoir for both WNV and SLEV. Many counties, nation-wide, have monitored deaths of birds known to be susceptible to WNV and have even tested them for the presence of WNV. Birds usually precede human cases by approximately two weeks (Kulasekera et al 2000; Mostashari et al 2003). This practice has been abandoned by many districts because it is expensive and labor intensive. A bird's extensive home ranges may not be a true representation of where WNV is a problem and the long-term effectiveness of a bird surveillance program remains uncertain due to natural selection of disease resistant birds (CDC, 2003). Therefore, surveillance of dead birds for WNV activity will be passive. TCPH will not test dead birds. If a citizen (or staff member) reports a dead bird sighting in the unincorporated area, we will record that information as a complaint and consider it as a possible surveillance location for an *ad hoc* mosquito trap. Should a dead bird sighting be reported in one of Tarrant County's municipalities, we will collect the information as a complaint, record it in a database and forward the information to the affected city.

Sentinel chicken flocks may be useful as a tool in WNV surveillance because they seroconvert (specific IgM antibodies in their blood that indicate WNV infection) at approximately the same time in the season as when human cases can be detected. Since WNV infections start with flu-like symptoms as do many other diseases, this may be helpful information to the medical community when making seasonal disease assessments within the region. However, this type of surveillance program should only be used in addition to other types of surveillance and after local transmission dynamics are well understood. It also may put workers at risk for infection while bleeding birds (CDC, 2003). TCPH has not implemented this type of surveillance..

\*Please note that CSTE case definitions may differ from doctor diagnoses. If a patient is diagnosed with WNV disease, an investigation by TCPH Epidemiology will follow to determine if this patient meets CSTE definition.

Manifestations of WNV infection are similar in horses to that of humans. According to the Texas Health and Safety Code (84.042), veterinarians are required to report disease in animals if it is on the list of reportable diseases for humans. Arboviral encephalitis including WNV and SLEV are also on the list of reportable diseases as listed in Section 97.3 (b) of the Texas Administrative Code, Communicable Diseases. (<http://www.dshs.state.tx.us/idcu/health/zoonosis/laws/report/>) Veterinary health professionals contact the Texas DSHS directly and thus TCPH is notified through the state when WNV infection has been confirmed in horses.

## **IV. Prevention and control**

All vector control personnel will be licensed by the Texas Department of Agriculture (TDA) for a non-commercial political license pertaining to vectors of diseases of public health concern within one year from their start date. Vector Control personnel will maintain all documents required by the Texas Department of Agriculture and Texas Commission of Environmental Quality (TCEQ) including the Texas Pollution Discharge Elimination System (TPEDS) General Permit. All vehicles will contain all necessary equipment and pertinent documentation for pesticide applications. All vehicles will be calibrated at least once per calendar year prior to the beginning of in-season WNV/SLEV surveillance. For a list of equipment and pertinent documentation see Appendix C. TCPH also has contracts in place for assistance with mosquito surveillance (trapping) and mosquito control activities (aerial and ground based) in cases where vector control staff are needed for other situations, or when vector control staff is overloaded.

### **A. Larval mosquitoes**

Larval surveillance is important in any integrated mosquito management program. Upon findings of a larval mosquito source, vector control technicians will bring a sample back to the lab. Upon returning to the lab, all larvae will be identified and the need for treatment will be assessed. Treatment of the larval source(s) will be determined on the estimated density and species (e.g. pest to humans or not). All treatments will be numbered in consecutive order of the time and date of treatment.

Reducing, eliminating, and treating mosquito breeding sites are some of the most important aspects of a mosquito control program. It controls mosquitoes before they emerge as host-seeking adults (only adult female mosquitoes are vectors for arboviral diseases). Additionally, pin-pointing sources that need to be treated are more efficient, effective, and safe than treating large areas with adulticide. Humans and terrestrial wildlife are less likely to be exposed to treated aquatic habitats of larval mosquitoes than those of terrestrial airborne adults.

Physical elimination and/or larviciding of mosquito breeding sites will be conducted by TCPH Vector Control personnel upon observation when possible. If sources are known to exist on private property, TCPH vector control technicians will notify the property owner of mosquito breeding activity

and request elimination and/or treatment of the breeding sources where possible. If the property owner does not eliminate/treat sources, TCPH may use applicable laws to seek legal remedy (Texas Health and Safety Code Chapters 341 and 343). See Appendix G for more information pertaining to larvicide.

*Gambusia affinis*, also known as mosquito fish, are natural endemic predators of aquatic stages of mosquitoes. They have the ability to continuously control mosquito larvae in a single source, are fairly easy to maintain, and are typically environmentally friendly. In partnership with the Tarrant Regional Water District (TRWD), vector control personnel will maintain *Gambusia* in four cement live wells near their Eagle Mountain Lake fish hatchery facility (See Appendix E for map). Vector control personnel will keep a stock tank of *Gambusia* at Tarrant County Public Health. Incorporated city personnel may request *Gambusia* at any time during business hours to be able to use for purposes of larval mosquito control within their municipalities. A record of the quantity of fish given and which city is asking for the fish will be maintained for stocking purposes.

## B. Adult mosquitoes

Surveillance of adult mosquitoes helps vector control technicians understand complex dynamics in mosquito ecology and the relationship between species. Different techniques of sampling will be utilized to assess different aspects of life within local adult mosquito populations. Since 2015 these have included gravid traps, CO<sub>2</sub>-baited light traps and BG Sentinel traps. The addition of New Jersey light traps and resting boxes are planned for 2019. Using different types of traps captures a wider variety of mosquito species. Specifically gravid traps target *Culex quinquefasciatus* and *Culex restuans*, whereas BG Sentinel traps target *Aedes aegypti* and *Aedes albopictus*.

Though larval elimination will be the main focus of mosquito control, it is impossible to know and eliminate every source of mosquito breeding. Mosquito borne illness can only be transmitted in the adult life stage of the mosquito vectors. TCPH continues to collect data to be able to establish infection rate and vector index threshold ranges that will help aid in adulticiding activity, but until these relationships are firmly established, Vector Control personnel or accepted contractors will continue to apply adulticide via ground-based, truck-mounted, Ultra Low Volume (ULV) sprayers in response to a mosquito sample that tests positive for WNV and/or SLEV. If an area continues to test positive for WNV and/or SLEV after it has been treated, TCPH will consider elevating this response to adulticiding multiple consecutive nights. Vector control personnel will also apply adulticide in response to a confirmed WNV human case after additional mosquito traps also test positive for WNV within a quarter of a mile radius around the location of the human case in accordance with CDC recommendations. As mentioned in the CHIKV/SLEV surveillance portion of this document, when large populations of adult *Aedes* are determined to be present on a suspected CHIKV/DENV patient's property, ULV or thermal fogging via backpack mounted/all-terrain vehicle (ATV) mounted sprayer will commence when appropriate. Tarrant County has a supplemental ground-based mosquito control contract with a commercial entity should additional assistance be required. Additionally, an aerial spray contract is in place for use during an arboviral related epidemic. See Appendix F for more information pertaining to adulticide and Tarrant County's phased response plan.

## **C. Resistance management**

Tarrant County Public Health uses integrated mosquito management techniques meant to minimize insect resistance. These include a combination of physical, chemical, biological and natural mosquito control techniques. Since there are limited classes of pesticides able to be used for control of adult mosquitoes (as opposed to the many options for larval control), TCPH will contract with an outside entity to conduct CDC approved resistance bioassays to determine adult mosquito resistance. If mosquitoes are determined to be resistant to a preferred adulticide, a secondary alternate adulticide will be utilized in future adulticiding activities. All possible contractors with TCPH will be notified of the adult mosquito conditions prior to treatment. It is recommended that municipalities consider testing their mosquito populations when these services are not available through TCPH.

## **V. Education, outreach, and media**

Educating the public is an essential aspect of Tarrant County's arboviral surveillance program. This includes giving the public tips on how to control mosquitoes in their homes and backyards, stating the risks and symptoms of these arboviral diseases and informing people on how to protect themselves from host-seeking female mosquitoes. To help educate the public, vector control personnel with the help of Medical Reserve Corps volunteers will attend local health fairs with tables including displays of larvae and adult mosquitoes, pamphlets and fliers that include information about mosquitoes, WNV/SLEV/CHIKV/DENV, and protection from mosquitoes, and answer general questions about these subjects. Vector control personnel and trained MRC volunteers will also be available for classes upon request to educate home owners associations (HOAs), elementary school classes, high-school classes and community outreach programs throughout the county. EHD and collaborating divisions will also be hosting an Arboviral Surveillance Program Kick-off meeting before the start of the WNV/SLEV season where vector control personnel will review the past year, review cumulative progress of the program and assess the program based on suggestions from municipalities. Vector control personnel will attend Texas Mosquito Control Association (TMCA) and American Mosquito Control Association (AMCA) meetings to be educated and to network with other vector control professionals when possible. Vector control personnel will work with our Public Information Officers (PIO)s to create original pamphlets, fliers, and other educational materials. These materials will be available for the public visiting Tarrant County Public Health as well as on the TCPH Website. TCPH's PIO handles official press releases, ground spray notifications, social media blogs, and other media involved items. Our PIO is also responsible for education campaigns, such as "Be Mosquito Free" and for the production of educational videos for the public about mosquito control.


## **VI. Mosquito complaints**

If mosquito breeding is suspected on any property within unincorporated Tarrant County, vector control personnel will conduct a thorough investigation of the suspect property. Complaints will be filed in the Tarrant County Larviciding Database. Complaints will be reviewed and prioritized in relation to severity of potential breeding sites (e.g. discarded tires, green swimming pools, observed containers holding stagnant water, etc.). Priority complaints will be investigated within 48 hours. After investigation vector control personnel will speak with the property owners to suggest remedy and contact complainant (if listed) to give an update of complaint status. All complaints will be numbered consecutively in the order they are received. Any required treatment will be referenced in the complaint. Vector control personnel will seek legal remedy when necessary (for list of applicable Texas Laws, see Appendix G).

## Appendix A: Trap Location

Vector control personnel have chosen trap locations based on sources of standing water, usually the first places which obtain water first, and the last to dry out. Temporary trap locations are chosen on a complaint basis, or when *Culex* sources are suspected by personnel. Trap locations were updated to maximize coverage of the unincorporated portions of the county in 2016. Cities choose their own locations. We recommend that cities choose permanent trap locations where sources are known to occur and/or where there is a historical presence of WNV/SLEV. They may also choose temporary trap locations on a complaint basis, or when scouting a new location. Good sources for *Culex quinquefasciatus* include catch basins, ditches with organically laden water, tires, and unkempt pools on abandoned properties.

Figure 1. Locations of in-season gravid trap locations in Tarrant County.



## **Appendix B: Internal notification of positive mosquito pools and trap location**

North Texas Regional Laboratory sends results of PCR testing for WNV/SLEV presence in mosquito pools from around the county to personnel within TCPH, including those in EHD. The Vector Control staff will record details of samples including the location coordinates, address, species and number of mosquitoes transferred to the tube, the total number of tested mosquitoes and the precinct in which the trap was placed. This information will be forwarded to the acting associate director or designee. The designee will notify appropriate contacts within Tarrant County, including the County Commissioners and the County Judge. Precinct contacts will also be notified for the purpose of placing appropriate signage.

## Appendix C: List of documents, emergency response, and other documentation

### General Documents

1. Certified applicator license
2. TDA pesticide application form
3. TPEDS general permit
4. Material Safety and Data Sheets (MSDS)s for chemicals used
5. Labels for chemicals used

### Emergency response

Occasionally during the loading or unloading of pesticide or in worst case, an accident, there may be spillage of the chemical. In the case of spillage action must be taken to contain and clean up the chemical. In addition, a person may have come in contact with the chemical.

If the chemical gets on someone, have the person leave the area immediately, wash quickly and thoroughly, change clothes and see a doctor if necessary (chemical is inhaled, swallowed, or comes in contact with highly susceptible areas such as the eyes).

Clear the area except for the cleanup staff. The cleanup staff must wear proper personal protective equipment and clothing.

When there is a spill, action is to be taken immediately to reduce the spill area and protect sensitive areas. Pet litter will be used as an absorbent to soak up the liquid. This material, along with soil that may have been impacted can be swept or scooped up and stored in a water proof container for proper disposal. A detergent and water can be used to clean-up the remainder of the residue if this appropriate for the surface but this water must also be collected for disposal.

### Personal protective equipment (PPE)

Any person involved in the transporting, handling, or dispensing must be wearing long pants, a long sleeved shirt, shoes, and socks. The label for the chemical will list required and recommended personal protective equipment (PPE) but at a minimum the follow items are to be available:

- Head covering
- Non-absorbent covering for clothes
- Chemical resistant gloves
- Eye goggles or face shield

- Respirator or some type of approved mouth covering
- Chemical resistant footwear

Though only the person handling the chemical is required to wear the prescribed PPE it is highly recommended that all personnel on the team have equipment available to them. Each person is to have a change of clothes available and there is to be backup equipment for disposable items such as gloves, respirator filters or mouth covering, or other items outlined by the label.

## Spill kit

The Spill Kit shall contain the following items:

- |                                      |                                  |
|--------------------------------------|----------------------------------|
| - At least five pounds of pet litter | - At least five gallons of water |
| - Broom                              | - Dust pan                       |
| - A container to collect waste       | - Detergent                      |
| - Chemical resistant gloves          | - Paper towels                   |
| - Eye/hand wash station              | - List of emergency contacts     |
| - First Aid Kit                      |                                  |

## Other documents

1. Trap data surveillance forms
2. Maintenance of equipment logs

Since County vehicles will be used for transporting the aduenticiding equipment the service schedule outlined by County Transportation will also be adhered to. The following maintenance checks will be conducted each day by staff prior to using the truck or trailer:

- |                               |
|-------------------------------|
| - Engine oil level            |
| - Amber light working         |
| - Horn                        |
| - Tire air pressure           |
| - All signal and break lights |
3. *Gambusia* log
    - Names of city personnel and approximate number of fish will be recorded by the vector control personnel each time fish are distributed.



Below is an example of a backyard checklist. It should be made available for uniform inspections of properties during backyard investigations.



Street Name: \_\_\_\_\_

### Mosquito Investigation Form - Backyard Checklist

CITY USE ONLY

**Investigation Instructions:**

- 1.) Inspect back and front yard using the Back Yard Check List.
- 2.) Set BG Sentinel trap in or near the yard. (May through November)
- 3.) Collect BG Sentinel trap the next afternoon.
- 4.) Fill out the Mosquito Surveillance form and deliver sample to TCPH Vector Control. 1101 S. Main St., Rm. 2300, Fort Worth, TX. 76104
- 5.) TCPH Vector Control will contact you with risk status and recommendations.
- 6.) If treatment is recommended and conducted, set a BG Sentinel trap after treatment.
- 7.) Deliver post - treatment BG Sentinel Trap to TCPH Vector Control for further evaluation.

Did you provide a Home Care Kit to Patient?	<input type="checkbox"/> Yes <input type="checkbox"/> No	Date: _____
Were there mosquito larvae present on the patient's property?	<input type="checkbox"/> Yes <input type="checkbox"/> No	
Were there adult mosquitoes present on the patient's property?	<input type="checkbox"/> Yes <input type="checkbox"/> No	
Does the patient have mosquitoes inside their house?	<input type="checkbox"/> Yes <input type="checkbox"/> No	

Checklist:		Observations / Actions / Recommendations
Bird Baths / Fountains	<input type="checkbox"/>	
Flower Pots	<input type="checkbox"/>	
Wheel Barrows	<input type="checkbox"/>	
Gutters	<input type="checkbox"/>	
Buckets	<input type="checkbox"/>	
Boats / Boat Covers	<input type="checkbox"/>	
Swimming Pool / Wading Pools	<input type="checkbox"/>	
Hot Tub	<input type="checkbox"/>	
Tires	<input type="checkbox"/>	
Animal Dishes / Troughs	<input type="checkbox"/>	
French Drains	<input type="checkbox"/>	
Child Play Stations	<input type="checkbox"/>	
Ornamental Ponds	<input type="checkbox"/>	
Tarp Covers	<input type="checkbox"/>	
Other Open Container Misc.	<input type="checkbox"/>	
Other Open Container Misc.	<input type="checkbox"/>	
Other Open Container Misc.	<input type="checkbox"/>	
Other Open Container Misc.	<input type="checkbox"/>	

Contacts: Nina Dacko (817) 321-4986 Shannon Solberg (817) 321-4989 Bethany Hambrick (817) 321-4971  
 Ivana Terry (817) 321-5364 Joe Carr (817) 321-5372



Below is an example of our Response Data Capture Form:

**Tarrant County Public Health**  
Accountability. Quality. Innovation.

  
A healthier community through leadership in health strategy

3 of 4

---

**Mosquito Investigation Form - Response Data Capture Form**

CITY USE ONLY

Did you conduct community education?:  Yes  No      Date: \_\_\_\_\_      How did you educate?: \_\_\_\_\_

Number of houses contacted?: \_\_\_\_\_      Number of houses you anticipated contacting?: \_\_\_\_\_

Did you apply any larvicide in response?:  Yes  No      Date/s: \_\_\_\_\_

What type of larvicide used?: \_\_\_\_\_      How applied?: \_\_\_\_\_

How many houses received larvicide treatment?: \_\_\_\_\_

Did you contract larvicide service with a company?:  Yes  No      What company?: \_\_\_\_\_

Did you apply barrier treatment?:  Yes  No      Date/s: \_\_\_\_\_

What kind of barrier treatment used?: \_\_\_\_\_      How applied?: \_\_\_\_\_

How many houses received barrier treatment?: \_\_\_\_\_

Did you contract barrier treatment with a company?:  Yes  No      What company?: \_\_\_\_\_

Did you apply adulticide treatment?:  Yes  No      Date/s: \_\_\_\_\_

What kind of adulticide used?: \_\_\_\_\_      How applied?: \_\_\_\_\_

How many houses received adulticide treatment?: \_\_\_\_\_

Did you contract adulticide treatment with a company?:  Yes  No      What company?: \_\_\_\_\_

Did you adulticide from the road?:  Yes  No      What kind of adulticide used?: \_\_\_\_\_

Did any houses refuse treatment?:  Yes  No      Number who refused larvicide treatment: \_\_\_\_\_

Number who refused barrier treatment: \_\_\_\_\_      Number who refused adulticiding: \_\_\_\_\_

Addresses of houses that refused treatment:	What type of treatment refused?:	Reason for refusal:

Contacts: Nina Dacko (817) 321-4986    Shannon Solberg (817) 321-4989    Bethany Hambrick (817) 321-4971  
Ivana Terry (817) 321-5364    Joe Carr (817) 321-5372

## Appendix E: Larvicides

### Microbial

#### Bacillus thuringiensis israelensis (Bti)

Bti is environmentally friendly in most situations because it is specific to insect pest species. This means that it will not harm beneficial insects that are natural predators of mosquitoes such as dragonfly nymphs, belostomatids and other aquatic invertebrates and/or fish. The mosquitoes must ingest this insecticide to work, so it is ineffective if a source is discovered when mosquito larvae are in the later part of their fourth instar when larvae stop eating to prepare for metamorphosis. It will also not be effective against pupae who do not eat. This bacterium produces a crystalline toxin that essentially destroys the larval midgut (digestive tract). Bti and Bs must be periodically reapplied.

#### Lysinibacillus sphaericus formerly Bacillus sphaericus (Bs)

Mode of action is similar to Bti.

### Spinosad

Spinosad is an insecticide that is based on the soil bacterium *Saccharopolyspora spinosa*. It has relatively low toxicity to birds and mammals, but is a broadrange pesticide. This means it may harm insects other than mosquitoes and so should not be used in benthic environments with an abundance of beneficial insects. Unlike Bti, it does not have to be ingested and should be effective against larvae of all instars and pupae. It alters the function of the nicotinic (acetylcholine) and GABA ( $\gamma$  aminobutyric acid) gated ion channels. This results in hyperexcitation of the muscles ultimately resulting in paralysis and death similar to the outcome of organophosphates.

### Chemical/Physical

#### Insect Growth Regulators (IGRs)

An insect growth regulator is a chemical which interrupts a portion of an insect's life cycle. Some of these may be as simple as a chitin inhibitor- meaning that it disrupts forming of the insect exoskeleton. There are also other types of IGRs that may behave similarly to endogenous or naturally occurring chemicals. As with most animals, many internal physiological functions are a result of the presence or absence of hormones. In insects, one of these functions is metamorphosis. When an insect is in its larval stage, a hormone known as juvenile hormone (JH), which is mostly order specific, is constantly present (although much less during the molting process). It is the absence of this hormone that causes the insect to enter into the metamorphic stage where it will transform into an adult insect. An insect growth regulator may be an analog, or similar molecules to JH, or any chemical that changes or

affects the growth and development of an insect. Therefore, in the presence of an insect growth regulator, the insect can never become an adult. This in turn disrupts the insect life cycle resulting in death. This should not affect the food web and may be used in a variety of environments. It is not recommended for larger bodies of water, or flowing water, but small places like septic systems, storm drains, catch basins, roadside ditches and the like. It does have to be periodically reapplied.

### **Monomolecular Films (MMF)**

A monomolecular film modifies the surface tension of the water. Mosquito larvae rest at the top of the water by placing their siphons at the top of the water where the natural surface tension keeps them afloat without activity. When the surface tension is modified, the mosquito larvae have to constantly swim to the top of the water to breathe. This results in exhaustion of the mosquito larvae (and pupae) and eventually death. This is not harmful to fish and other wildlife. Since the mode of action of this substance is physical, mosquitoes cannot build up resistance. It does require reapplication every few weeks. Since MMFs are self-spreading, if applied correctly, they have great coverage and may be used in larger environments like rice fields, swamps and wetlands.

### **Oils**

Oils are insecticides that spread over the top of the water to create a barrier. Mosquito larvae and pupae cannot push their breathing apparatus through this oil resulting in suffocation. This is one of the quickest ways to eradicate mosquito larvae and is another insecticide mosquitoes cannot become resistant to because of the physical mode of action. These products should not be used in the presence of other aquatic wildlife because it can affect the dissolved oxygen content and also contains petroleum distillate.

## Appendix F: *Gambusia affinis*

Our *Gambusia* program stocked live wells are located at the Tarrant Regional Water District Eagle Mountain fish hatchery, located at 8665 Eagle Mountain Circle, Fort Worth TX, 76135 in the northwestern portion of Tarrant County. Vector Control personnel will obtain badges to be able to enter Tarrant Regional Water District Facilities.

Figure 2. Aerial picture of live wells for storage of *Gambusia affinis*





Figure 3. Location of TRWD fish hatchery in northwest Tarrant County



## Appendix G: Adulticides

### Chemical

#### Pyrethroids

Pyrethroids are a class of pesticides which are derived from botanicals found in certain species of flowers in the genus *Chrysanthemum*. They are non-persistent in the environment and are broken down in a short amount of time by UV light (sunlight). These insecticides may be broad range (meaning they can affect many types of insects). To avoid beneficial pollinators such as bees and butterflies, and to optimize the number of adult mosquitoes killed, these insecticides are typically sprayed at night when mosquitoes are most active and pollinators are resting. These chemicals are also applied with ultra-low volume (ULV) equipment, meaning very little of the insecticide is needed for treatment. Tiny droplets of insecticide must come in contact with the adult mosquito for it to be effective. Pyrethroids affect the gated ion channels of the insect's nervous system in such a way to where the muscles become over stimulated, which eventually leads to paralysis and death. Bodies of water should be avoided when using these chemicals because they can be toxic to other benthic wildlife including fish and beneficial insects.

#### Organophosphates

Organophosphate pesticides should only be used for adult mosquito control in instances where the local adult population shows resistance to pyrethroid insecticides. Organophosphates, in large doses, may be toxic to other wildlife and humans, birds and other mammals. The doses required for mosquito control are well below these levels (also applied with ULV equipment). Most organophosphates also break down quickly in sunlight. These pesticides affect acetylcholinesterase which break down acetylcholine, a neurotransmitter that controls muscle function. Inability to break down this neurotransmitter leads to overstimulation of the muscles, and eventually paralysis and death.

## Appendix H. Phased response Guidelines

**Table 1 Tarrant County Public Health Guidelines for Phased Response to WNV/SLEV Surveillance**

The following recommendations are intended to guide mosquito control programs and may include other applicable community procedures. All actions are subject to change without notice due to organizational priorities, weather or other unforeseen circumstances.

Risk category	Probability of human outbreak	Public Health Threshold	Recommended response
0	None	No evidence of mosquito or viral activity	<p><b>Surveillance</b></p> <ul style="list-style-type: none"> <li>Develop response plan</li> </ul> <p><b>Information/Education</b></p> <ul style="list-style-type: none"> <li>Initiate community outreach and public education programs</li> <li>Conduct audience research to develop/target education &amp; community involvement</li> </ul> <p><b>Control Measures</b></p> <ul style="list-style-type: none"> <li>Contact community partners</li> <li>Secure necessary control resources to enable emergency response</li> </ul>
1	Low	Normal mosquito activity with little or no evidence of viral activity	<p><b>Surveillance</b></p> <ul style="list-style-type: none"> <li>Routine monitoring of public health threats</li> <li>Monitor larval and adult mosquito bionomics (trap surveillance)</li> <li>Identify disease-causing agents in mosquito samples through laboratory testing;</li> </ul> <p><b>Information/Education</b></p> <ul style="list-style-type: none"> <li>Public education and community outreach programs focused on risk potential, personal protection and residential source reduction</li> </ul> <p><b>Control Measures</b></p> <ul style="list-style-type: none"> <li>Source reduction through physical and environmental measures</li> <li>Biological – mosquito fish</li> <li>Larvicides (surface oils, biorational bacterial products and insect growth regulators)</li> </ul>
2	Moderate	Virus detected in mosquito samples	<p><b>Surveillance</b></p> <ul style="list-style-type: none"> <li>Increase mosquito surveillance in areas of positive traps</li> </ul> <p><b>Information/Education</b></p> <ul style="list-style-type: none"> <li>Public health advisory released</li> <li>Advise the public and emphasize source reduction, personal protection and disease symptoms</li> </ul> <p><b>Control Measures</b></p> <ul style="list-style-type: none"> <li>Larviciding of breeding sites around positive trap sites</li> </ul>

			<ul style="list-style-type: none"> <li>• Consider ground-based ULV adulticide application around positive trap location, with continued mosquito surveillance</li> <li>• Consider aerial application if detection of virus is widespread and when infection rates in mosquitoes are high and increasing, even in the absence of human cases</li> </ul>
3	High	Virus detected in multiple mosquito samples from different times and locations and confirmed human cases	<p><b>Surveillance</b></p> <ul style="list-style-type: none"> <li>• Increase surveillance activities in adjacent areas where spread of virus is likely</li> </ul> <p><b>Information/Education</b></p> <ul style="list-style-type: none"> <li>• Public health warning released</li> <li>• Publicize vector control measures within the target communities</li> <li>• Warn the general public of the probability of disease and provide guidance</li> </ul> <p><b>Control Measures</b></p> <ul style="list-style-type: none"> <li>• Ground-based ULV adulticide application of area around confirmed human case(s) after mosquito trapping confirms the presence of disease</li> <li>• Ground-based ULV adulticide application of positive mosquito sites and expanded areas around sites</li> <li>• Consider aerial treatment(s)</li> </ul>
4	Outbreak in progress	Multiple human cases confirmed and detection of increased or continued viral mosquito activity	<p><b>Surveillance</b></p> <ul style="list-style-type: none"> <li>• Continue mosquito surveillance in areas of confirmed human cases</li> </ul> <p><b>Information/Education</b></p> <ul style="list-style-type: none"> <li>• Public health emergency considered</li> <li>• Declaration of public health emergency/distribution of emergency alerts</li> </ul> <p><b>Control Measures</b></p> <ul style="list-style-type: none"> <li>• Ground-based ULV adulticide application in areas of clustered human cases</li> <li>• Recommend aerial adulticide applications in targeted zones</li> </ul>

Tarrant County mosquito response guidelines were adapted from the Centers for Disease Control and Prevention report "Epidemic/Epizootic West Nile Virus in the United States: Guidelines for Surveillance, Prevention and Control, 2013"

**Table 2 Tarrant County Public Health Guidelines for Phased Response to human cases of chikungunya, dengue and Zika**

Risk Category	Definition	Jurisdictional Action Steps
Preparation	Vector Present or Possible in Jurisdiction	<p><b>Vector Control Preparation:</b></p> <ul style="list-style-type: none"> <li>• Prepare for surveillance of <i>Aedes aegypti</i> and <i>Aedes albopictus</i> by deploying BG Sentinel traps.</li> <li>• Recommend municipalities to enforce and encourage community clean-up efforts.</li> <li>• Disseminate public education materials to municipalities &amp; county residents.</li> <li>• Update plans for mosquito reduction tactics around travel-associated cases.</li> <li>• Update appropriate contracts for Zika-type response including localized adulticiding and larviciding.</li> <li>• Update permission forms for entrance into private residences and standardized response backyard checklists.</li> <li>• Communicate Tarrant County response plan with each municipality individually.</li> <li>• Reiterate response plan during Arthropod-borne diseases kick-off meeting.</li> <li>• Increase vector personnel and expand abilities to respond to surveillance and control needs.</li> <li>• Plan for Resistance Testing</li> <li>• Research new methods of <i>Aedes</i> spp. control</li> </ul>
Category 1	<p><b>Mosquito Season</b> (<i>Aedes aegypti</i> or <i>Aedes albopictus</i> mosquito biting activity)</p>	<p><b>Continue preparation category activities in addition to the following steps:</b></p> <p><b>Vector Control:</b></p> <ul style="list-style-type: none"> <li>• Monitor <i>Aedes aegypti</i> and <i>Aedes albopictus</i> populations.</li> <li>• Identify potential <i>Aedes</i> spp breeding sources and concentrate on eliminating the sources of the mosquitoes, continue public education efforts and encourage community cleanup.</li> <li>• Consider distributing Zika care kits supplied by Tarrant County Public Health to all suspect or confirmed Zika/CHIK/DEN patients.</li> <li>• After identification of a <b>suspected</b> travel-associated human case outside the viremic phase, consider educating the patient about mosquito breeding sources and/or showing them how to perform backyard inspections.</li> </ul>

		<ul style="list-style-type: none"> <li>• After identification of a <b>suspected</b> travel-associated human case within the viremic phase of disease:</li> <li>• Consider inspecting the patient’s yard for mosquito larval sources/activity.</li> <li>• Consider gaining entry into property and utilize permission form.</li> <li>• Upon entry to the private property, record any larval or adult activity on the property.</li> <li>• Set a BG trap within the mosquito season to determine vector abundance and help determine risk.</li> <li>• If mosquito activity is observed, consider larviciding, pupiciding and/or adulticiding the property.</li> <li>• Consider residual barrier treatments for adulticiding responses</li> <li>• Thermal fogging may be advantageous in serious problem areas with high adult activity (such as tire piles or dumping areas).</li> <li>• Set a BG Sentinel trap post treatment to monitor and assess effectiveness of vector control activities.</li> <li>• After identification of a <b>confirmed</b> travel-associated human case outside of the viremic phase of disease:</li> <li>• Source reduction, larvicide and personal protective information should be supplied to houses up to a 150 meter radius within the block location.</li> <li>• After identification of a <b>confirmed</b> travel-associated human case within the viremic phase of disease, consider taking the same measures as one would in the suspected travel-associated human case with adjacent properties to the patient property.</li> <li>• Source reduction, larvicide and personal protective information should be supplied to houses up to a 150 meter radius where mosquito activity is observed around a travel-associated case.</li> <li>• Intensify community cleanup up to a 150 meter radius around suspected case.</li> <li>• Consider utilizing autocidal gravid trapping strategies for additional adult control.</li> <li>• Continue to monitor information distributed by the CDC for the most up to date information.</li> </ul>
--	--	--

<p><b>Category 2</b></p>	<p><b>Confirmed Local Transmission</b> (single case, or cases clustered in a single household/community in a county or jurisdiction)</p>	<p><b><i>Continue category 1 activities, in addition to the following steps.</i></b></p> <p><b>Vector Control:</b></p> <ul style="list-style-type: none"> <li>• Expand larvicide, pupicide and adulticiding activities up to 150 meter radius around initial case(s) in addition to supplying source reduction, larvicide and personal protective information and intensified community cleanup.</li> <li>• Monitor <i>Aedes aegypti</i> and <i>Aedes albopictus</i> activity around suspected local transmission with BG Sentinel traps.</li> <li>• Continue to assess effectiveness of vector control activities.</li> </ul>
<p><b>Category 3</b></p>	<p><b>Widespread Local Transmission</b> (multiple locations within a county or jurisdiction)</p>	<p><b><i>Continue category 2 activities, in addition to the following step.</i></b></p> <p><b>Vector Control:</b></p> <ul style="list-style-type: none"> <li>• Consider expanding areas and utilizing ground based and/or aerial control adulticiding/larviciding activities.</li> <li>• Intensify community cleanup with messages on media.</li> <li>• Intensify public education regarding the need of mosquito control and personal protective measures.</li> <li>• Continue to assess effectiveness of vector control activities</li> </ul>
<p><b>Category 4</b></p>	<p><b>Local Transmission in Multiple Counties</b></p>	<p><b><i>Continue category 3 activities, in addition to the following step.</i></b></p> <ul style="list-style-type: none"> <li>• Expand response activities regionally or state-wide</li> </ul>

## **Appendix I: Laws pertaining to vector control**

### **Texas Health and Safety Code**

#### **Title 5, Subtitle A, Chapter 341, Subchapter B**

.011- Defines public health nuisance

.012- Requires abatement of nuisance

.019- Public officials can treat an abandoned or uninhabited property with mosquito breeding sites

#### **Title 5, Subtitle A, Chapter 343, Subchapter B**

.011- maintain property so as not to attract mosquitoes, rodents, vermin or disease-carrying pests

.012- States penalty from \$50-\$200 if nuisance is not abated within 30 days of notice; if previously convicted \$200-\$1000 and/or jail time up to six months. Each day the violation occurs is a separate offense.

#### **Title 5, Subtitle A, Chapter 341, Subchapter C**

.024- Authority to enter premises when presenting proper government identification

Federal Insecticide, Fungicide and Rodenticide Act (FIFRA) (entirety- bound by pesticide license)

Clean Water Act, Section 402 and Chapter 26 of the Texas Water Code- pertains to a general permit to discharge biological pesticides and chemical pesticides that leave a residue in water in or near waters of the US.

## Glossary of acronyms

AAM- American Academy of Microbiology

AMCA- American Mosquito Control Association

ATV- All-Terrain Vehicle

Bs- *Bacillus sphaericus*

Bti- *Bacillus thuringiensis israelensis*

CDC- Centers for Disease Control and Prevention

CHIK- Chikungunya fever

CHIKV- Chikungunya virus

CSTE- Council of State and Territorial Epidemiologists

DEET- N, N, diethyl-*meta*-toluamide

DENV- Dengue virus

DSHS- Department of State Health Services

EHD- Environmental Health Division

GABA- Gamma AminoButyric Acid

HIPAA- Health Insurance Portability and Accountability

Act

HOA- Home Owners Association

IGR- Insect Growth Regulator

JH- Juvenile Hormone

MMF- MonoMolecular Film

NTRL- North Texas Regional Laboratory

PAHO- Pan American Health Organization

PIO- public information officer

PPE- Personal Protective Equipment

RT-PCR- Reverse Transcriptase Polymerase Chain

Reaction

SLEV- St. Louis encephalitis virus

TCEQ- Texas Commission on Environmental Quality

TCPH- Tarrant County Public Health

TDA- Texas Department of Agriculture

TMCA- Texas Mosquito Control Association

TPEDS- Texas Pollution Elimination Discharge System

TRWD- Tarrant Regional Water District

ULV- Ultra Low Volume

WNV- West Nile virus

ZIKV- Zika virus

## Works cited

- American Academy of Microbiology. West Nile Virus: FAQ. Washington DC. AAM. 2013.
- Ward, M. P. Spread of equine West Nile virus encephalomyelitis during the 2002 Texas epidemic. *Am. J. Trop. Med. Hyg.*, 74(6) 2006, pp.1090-1095
- Rasmussen, S. A., D. J. Jamieson, M. A. Hoenien, L. R. Petereson. Zika Virus and Birth Defects- Reviewing the Evidence for Causality. *N. Engl. J. Med.* 2016. 374 pp. 1981-1987
- Strickman, D. 1988. Rate of oviposition by *Culex quinquefasciatus* in San Antonio, Texas, during three years. *J. Am. Mosq. Con. Ass.* 1988. 3(3), pp. 339-344
- Pan American Health Organization, Preparedness and Response for Chikungunya virus: Introduction into the Americas. Washington DC. PAHO. 2011.
- World Health Organization. Dengue: Guidelines for Diagnoses, Prevention, Treatment and Control. Geneva, Switzerland. WHO. 2009.
- Richards, S. L., S. I. Anderson and B. W. Alto, Vector Competence of *Aedes aegypti* and *Aedes albopictus* (Diptera: Culicidae) for Dengue Virus in the Florida Keys. *J. Med. Entomol.* 2012. 49 (4), pp 942-946
- Rodriguez-Tan, R. S. and M. R. Weir. Dengue: a review. *Tx. Med.* 1998. 94 (10) pp. 53-9.
- Mostashari, F; M. Kulldorf; J. J. Hartman; J. R. Miller; V. Kulasekera. Dead bird clusters as an early warning system for West Nile virus activity. *Emerg. Inf. Dis.*, 9 (6) 2003. pp. 641-646
- Centers for Disease Control and Prevention. West Nile virus in the United States: Guidelines for surveillance, prevention, and control. 4<sup>th</sup>Revision. CDC. 2013.
- ArboNET, Centers for Disease Control and Prevention: Arboviral Diseases Branch. West Nile virus human cases reported to CDC by state 2002-2012.  
[http://www.cdc.gov/westnile/resources/pdfs/cummulative/99\\_2012\\_cummulativeHumanCases.pdf](http://www.cdc.gov/westnile/resources/pdfs/cummulative/99_2012_cummulativeHumanCases.pdf)
- Department of State Health Services. Veterinary Responsibilities in Disease Reporting. Updated 5/2013.  
<http://www.dshs.state.tx.us/idcu/health/zoonosis/laws/report/>